



ELSEVIER

Contents lists available at ScienceDirect

European Journal of Pharmacology

journal homepage: www.elsevier.com/locate/ejphar

Full length article

Gamma-linolenic acid, Dihommo-gamma linolenic, Eicosanoids and Inflammatory Processes

Susan Sergeant^a, Elaheh Rahbar^b, Floyd H. Chilton^{c,*}^a Department of Biochemistry; Wake Forest School of Medicine, One Medical Center Blvd, Winston-Salem, NC 27157, USA^b Department of Biomedical Engineering; Wake Forest School of Medicine, One Medical Center Blvd, Winston-Salem, NC 27157, USA^c Department of Physiology/Pharmacology, Wake Forest School of Medicine, One Medical Center Blvd, Winston-Salem, NC 27157, USA

ARTICLE INFO

Article history:

Received 23 February 2016

Received in revised form

25 March 2016

Accepted 11 April 2016

Keywords:

Gamma-linolenic acid

Dihommo gamma-linolenic acid

Arachidonic acid

Eicosanoid

Inflammation

ABSTRACT

Gamma-linolenic acid (GLA, 18:3n-6) is an omega-6 (n-6), 18 carbon (18C-) polyunsaturated fatty acid (PUFA) found in human milk and several botanical seed oils and is typically consumed as part of a dietary supplement. While there have been numerous *in vitro* and *in vivo* animal models which illustrate that GLA-supplemented diets attenuate inflammatory responses, clinical studies utilizing GLA or GLA in combination with omega-3 (n-3) PUFAs have been much less conclusive. A central premise of this review is that there are critical metabolic and genetic factors that affect the conversion of GLA to dihommo-gamma linolenic acid (DGLA, 20:3n-6) and arachidonic acid (AA, 20:4n-6), which consequently affects the balance of DGLA- and AA- derived metabolites. As a result, these factors impact the clinical effectiveness of GLA or GLA/(n-3) PUFA supplementations in treating inflammatory conditions. Specifically, these factors include: 1) the capacity for different human cells and tissues to convert GLA to DGLA and AA and to metabolize DGLA and AA to bioactive metabolites; 2) the opposing effects of DGLA and AA metabolites on inflammatory processes and diseases; and 3) the impact of genetic variations within the fatty acid desaturase (*FADS*) gene cluster, in particular, on AA/DGLA ratios and bioactive metabolites. We postulate that these factors influence the heterogeneity of results observed in GLA supplement-based clinical trials and suggest that "one-size fits all" approaches utilizing PUFA-based supplements may no longer be appropriate for the prevention and treatment of complex human diseases.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Gamma-linolenic acid (GLA, 18:3n-6) is an omega-6 (n-6), 18 carbon (18C) polyunsaturated fatty acid (PUFA) found in human milk and several botanical seed oils (borage [\sim 21% GLA], blackcurrant [\sim 17%GLA] and evening primrose [\sim 9%GLA]), and is typically consumed as a part of a dietary supplement. The scientific literature examining the clinical effects of GLA-containing supplements is both complex and confusing. While there have been numerous *in vitro* and *in vivo* animal models illustrating that GLA-supplemented diets attenuate various inflammatory responses, the clinical literature has been less conclusive (for a review, see (Fan and Chapkin, 1998)). The introduction of GLA supplementation strategies to achieve symptomatic relief of atopic dermatitis/eczema was historically preceded by the use of relatively large daily doses of oral linoleic acid (LA, 18:2n-6) containing oils. This

was based on the premise that patients with atopic dermatitis/eczema had hallmark cutaneous signs of essential fatty acid deficiency and an impairment in PUFA biosynthesis at an early desaturase step (*FADS2*; Δ -6 desaturase) (Burr and Burr, 1929; Burr et al., 1932; Horrobin, 2000). It was hypothesized that GLA supplements could restore needed PUFAs and mitigate the disease.

Numerous studies primarily carried out in the 1980s and 1990s demonstrated that GLA-enriched botanical oils (evening primrose, borage, blackcurrant seed, and fungal-derived) had the capacity to relieve the signs and symptoms of several chronic inflammatory diseases, including rheumatoid arthritis (RA) and atopic dermatitis (Andreassi et al., 1997; Foolad et al., 2013; Kunkel et al., 1981; Leventhal et al., 1994, 1993; Lovell et al., 1981; Morse et al., 1989; Tate et al., 1989; Zurier et al., 1996). However, several more recent reviews and meta-analyses have questioned these earlier studies and raised doubts about the clinical effectiveness of GLA-enriched supplements particularly in the context of atopic dermatitis and RA (Bamford et al., 2013; Belch and Hill, 2000; Kitz et al., 2006; Macfarlane et al., 2011; Van Gool et al., 2004) (Table 1). A variety of issues complicate these studies including the fact that many of the

* Corresponding author.

E-mail addresses: ssergean@wakehealth.edu (S. Sergeant), rahbar@wakehealth.edu (E. Rahbar), schilton@wakehealth.edu (F.H. Chilton).

trials have: 1) relatively low subject numbers; 2) less than ideal study designs (e.g. the absence of washout period in cross-over design trials); 3) variations in the types of GLA supplements and how they are administered (e.g. dose, duration); and 4) differences in selection/inclusion criteria (e.g. population demographics and disease states) (Foster et al., 2010; Van Gool et al., 2004).

Several studies have also investigated the effects of GLA when given in combination with botanical or marine omega-3 (n-3) enriched PUFA supplements. Enteral diets enriched with marine oils containing (n-3) LC-PUFAs (i.e. eicosapentaenoic acid [EPA, 20:5n-3] and docosahexaenoic acid [DHA, 22:6n-3]) and GLA have been shown to reduce cytokine production and neutrophil recruitment into the lung resulting in fewer days on ventilation and shorter stays in the intensive care unit in patients with acute lung injury or acute respiratory distress syndrome (Gadek et al., 1999; Pontes-Arruda et al., 2006; Singer et al., 2006). Importantly, these dietary combinations of GLA and (n-3) LC-PUFAs were also shown to reduce both morbidity and mortality of critically ill patients (Gadek et al., 1999; Li et al., 2015; Pontes-Arruda et al., 2006; Singer et al., 2006). However, as with the studies of GLA alone, the results combining GLA and (n-3) LC-PUFAs have not been reproducible. Other clinical studies, such as the OMEGA trial, did not show a benefit of these GLA/(n-3) LC-PUFA combinations on patient outcomes (Rice et al., 2011).

Supplementation strategies providing GLA together with (n-3) LC-PUFAs (i.e. EPA and DHA) have also been utilized in patients with atopic asthma (Surette et al., 2003a, 2003b, 2008) and have been shown to block *ex vivo* synthesis of leukotrienes from whole blood and isolated neutrophils. Importantly when provided as an emulsion, daily consumption of these combinations was associated with an improved quality of life in asthma patients and a decreased reliance on rescue medication (Surette et al., 2008). These results compared favorably with quality of life scores obtained in mild asthmatics treated with montelukast or zafirlukast (Riccioni et al., 2004). However, to our knowledge, no randomized, placebo-controlled trials have been conducted to investigate the effect of these combinations on the improved quality of life or other relevant clinical outcomes in asthma patients.

Alternatively, botanical oil combinations (e.g. borage and echium oils) containing GLA, the (n-3) 18C-PUFAs, alpha-linolenic acid (ALA, 18:3n-3) and stearidonic acid (SDA, 18:4n-3), have been shown to reduce leukotriene generation and forced expiratory volume in mild asthmatics (Arm et al., 2013; Kazani et al., 2014), improve glucose tolerance in insulin-resistant monkeys (Kavanagh et al., 2013) and reduce total and LDL cholesterol levels in patients with diabetes and metabolic syndrome (Lee et al., 2014). These botanical oil studies, however, have yet to be replicated in larger human clinical trials.

Together, these data indicate that the outcomes of clinical studies utilizing GLA supplementation, alone or in combination with other fatty acid-based supplements, while promising are highly inconsistent. These observations raise serious questions about our current understanding of the highly complex and dynamic nature of PUFA metabolism. More recent studies suggest that there are important metabolic and genetic factors within the human host that significantly impact the study of GLA or GLA/(n-3) PUFAs combinations and reveal that a "one size fits all" model of supplementation may not be appropriate. Further, these studies suggest that it may be necessary to better understand key metabolic and genetic issues regarding GLA metabolism before GLA-enriched supplements can be effectively used to address human disease. This review will focus on potential key metabolic and genetic factors that may impact the use and clinical effectiveness of GLA or GLA/(n-3) PUFA combinations.

2. Polyunsaturated fatty acid metabolism

2.1. Long chain polyunsaturated fatty acid biosynthesis

In mammals, (n-6) and (n-3), long chain (> 20 carbons, LC) PUFAs such as arachidonic acid (AA, 20:4n-6), dihomo-gamma linolenic acid (DGLA, 20:3n-6), EPA (20:5n-3) and DHA (22:6n-3) can be synthesized from their respective precursors, (n-6) and (n-3) 18C-PUFAs such as LA, GLA, ALA, and SDA. The PUFA pathways and attendant enzymes are illustrated in Fig. 1. Biologically important (n-6) LC-PUFAs, DGLA and AA can be synthesized from LA using either two (one desaturation and one elongation step) or three (two desaturation and one elongation step) enzymatic steps, respectively (Sprecher, 1981). The desaturation reactions have long been recognized as the rate-limiting steps in this pathway (Bernert and Sprecher, 1975) and the enzymes that catalyze these reactions are encoded by the fatty acid desaturase 1 and 2 (i.e. *FADS1* and *FADS2*) genes located on chromosome 11. This region is commonly referred to as the *FADS* cluster (11Q12.2-q13.1) (Glaser et al., 2011). These same enzymes are responsible for the rate-limiting steps in the conversion (n-3) 18C-PUFAs (ALA and SDA) to (n-3) long chain-PUFAs including EPA. Recent studies suggest that the efficiency of several steps in the pathway, in particular the desaturase steps, is highly impacted by genetic variations within the *FADS* cluster (Chilton et al., 2014; Eaton, 1992). The potential impact of these genetic variations on GLA, DGLA and AA levels and their respective ratios will be discussed in greater detail in Section 4. In addition, small quantities of LC-PUFAs can be obtained directly from the diet (Fig. 1). Dietary AA is obtained primarily from animal products such as, organ meats, eggs, poultry, and fish, whereas dietary EPA and DHA are found primarily in seafood, particularly cold-water fish (Hibbeln et al., 2006).

GLA enters the (n-6) pathway distal to the *FADS2* enzymatic step and is efficiently converted to DGLA by an enzymatic activity, encoded for by a gene known as *ELOVL5*, in a wide range of cells (including several inflammatory cells) and tissues. Because of its rapid conversion, GLA is found in low levels in circulating lipids, cells or tissues. In contrast to GLA, the *ELOVL5* product, DGLA is readily measured in circulating lipids and most cells, and levels of DGLA are consistently elevated after GLA supplementation (Chilton-Lopez et al., 1996; Johnson et al., 1997; Lee et al., 2014; Simon et al., 2014) (Table 2). Once formed, DGLA can be incorporated into cellular glycerolipids (primarily phospholipids). Upon cell activation, DGLA can be released as a free fatty acid by phospholipase A₂(s) and enzymatically converted to several metabolites with predominantly anti-inflammatory properties. These pathways are described in further detail in Section 3.3 below.

2.2. Metabolism of PUFAs to Lipid Mediators that Impact the Immune Response

DGLA and its metabolites have long been recognized to have potent inhibitory effects on platelet aggregation and inflammation. The impact of DGLA on platelet aggregation was first recognized in the early 1970s (Willis et al., 1974a), and Lagarde and colleagues showed that ten times more collagen and twice as much thrombin were necessary to obtain aggregation when platelets and endothelial cells were pretreated with DGLA as compared to untreated platelets and endothelial cells (Lagarde et al., 1981). Interestingly, DGLA was much more potent than EPA in inhibiting platelet aggregation.

The anti-inflammatory effects of DGLA have been attributed to both i) the anti-inflammatory properties of DGLA-derived metabolites and ii) the ability of DGLA to compete with AA in the synthesis of pro-inflammatory AA products (Billah et al., 1985; Chilton-Lopez et al., 1996; Iversen et al., 1991, 1992; Vanderhoek




Fig. 1. Illustration of PUFA pathway.

et al., 1980) (Table 3). The synthesis of DGLA oxygenated metabolites and their impact is discussed in detail below in Section 3.3.

Somewhat paradoxically from an inflammation perspective, AA can also be synthesized from DGLA utilizing an enzymatic activity originally known as the Δ -5 desaturase. As shown in Fig. 1, this activity is encoded for by *FADS1* within the *FADS* cluster. AA and its metabolic products have long been known to play important roles in immunity and inflammation (Boyce, 2008; Calder, 2013; Schmitz and Ecker, 2008; Simopoulos, 2008) via their ability to impact normal and pathophysiologic responses through the conversion of AA to potent eicosanoid products (including prostaglandins [PG], thromboxanes [TX], leukotrienes [LT] and lipoxins). Additionally, AA and its oxidized products can regulate transcription and consequently a wide range of cellular activities via cellular and nuclear receptors (such NF- κ B, PPAR and SREBP-1c (Berger et al., 2006; Caputo et al., 2011; Chinetti et al., 2000; Deckelbaum et al., 2006; Jump and Clarke, 1999; Jung et al., 2012; Schmitz and Ecker, 2008; Soberman and Christmas, 2003; Vanden Heuvel, 2012), thereby modulating the expression of numerous genes that impact immune responses. Therefore, dietary supplementation with GLA has the capacity to both increase levels of both DGLA, which can lead to several anti-inflammatory metabolites, and AA, whose metabolic products generally promote inflammation.

3. Factors that determine the balance of pro- and anti-inflammatory PUFAs and PUFA metabolites after GLA

supplementation

3.1. Differential metabolism of GLA to DGLA and AA in human cells and tissues

Since metabolites of DGLA have predominantly anti-inflammatory effects and AA products generally enhance inflammation, it stands to reason that the balance of AA to DGLA (i.e. the ratio of AA/DGLA) in circulation, cells and tissues is a critical factor that impacts inflammatory processes. Several factors determine the levels of AA and DGLA and thus the ratio of AA metabolites and DGLA metabolites within cells and tissues. One is the differential capacities of cells or tissues to elongate GLA to DGLA and then to further desaturate it to AA. Differential expression of enzymatic activities is observed when comparing GLA metabolism within an inflammatory cell, such as the human neutrophil, and within a tissue bed or organ, such as the human liver. Both *in vitro* and *in vivo* studies demonstrate that human neutrophils contain the elongase (*ELOVL5*) but not the Δ -5 desaturase (i.e. *FADS1*) activity. In addition to human neutrophils, skin, murine peritoneal macrophages and platelets also appear to have high *ELOVL5* elongase activity relative to *FADS1* (Δ -5) desaturase activity (Chapkin and Coble, 1991; Chapkin et al., 1988b; Chapkin and Ziboh, 1984; de Bravo et al., 1985; Navarette et al., 1992; Ziboh et al., 2000). In contrast, several other tissues including liver, kidney, testes, brain and intestine appear to contain both activities (Bernert and Sprecher, 2006; Blond and Bézard, 1991; Hurtado de Catalfo et al., 1992; Irazu et al., 1993; Luthria and Sprecher, 1994; Pawlosky et al., 1994).

The PUFA enzymatic portfolio of human neutrophils

Table 2
Effect of GLA supplementation on the DGLA content of immune cells.

Reference	design	supplement	dose	Duration	cell type tested	Biochemical outcomes			Unit
						[DGLA]-pre	[DGLA]-post	p value	
Fletcher and Ziboh (1990)	<i>In vivo</i> fatty acid supplementation	borage oil	3% oil (by weight) in diet	12wk	guinea pig neutrophil	0.27	2.41	p ^a	mg /100 mg total phospholipid
Ziboh and Fletcher (1992)	<i>In vivo</i> fatty acid supplementation	borage oil	0.48 g GLA/d	6wk	human neutrophil	1.35	1.96	<0.05	mg /100 mg total phospholipid
Ziboh and Fletcher (1992)	<i>In vivo</i> fatty acid supplementation	blackcurrant oil	0.48 g GLA/d	6wk	human neutrophil	1.35	1.95	<0.05	mg /100 mg total phospholipid
Chilton-Lopez et al. (1996)	<i>In vitro</i> fatty acid incorporation	albumin-conjugated GLA	200nmol	24 h	human neutrophil	0.4	1.0	p ^a	nmol
Chilton-Lopez et al. (1996)	<i>In vivo</i> fatty acid supplementation	borage oil	3 g GLA/d	21d	human neutrophil	0.45	0.80	<0.05	nmol/million cells
Johnson et al. (1997)	<i>In vivo</i> fatty acid supplementation	borage oil + controlled diet	1.5 g GLA/d	21d	human neutrophil	0.20	0.20	ns ^b	nmol/5 million cells
Chapkin and Coble (1991)	<i>In vitro</i> fatty acid incorporation	albumin-conjugated radiolabeled GLA	tracer amounts	21d	human neutrophil	0.15	0.27	<0.05	nmol/5 million cells
				21d	human neutrophil	0.13	0.35	<0.05	nmol/5 million cells
Ziboh et al. (2004)	<i>In vivo</i> fatty acid supplementation	borage oil	2 g GLA/d	3hr	murine peritoneal macrophage	0	84	p ^a	% conversion to radiolabel product
				12mo	human neutrophil	0.7	1.4	<0.05	mg /100 mg total phospholipid

^a p value not specified.

^b ns, not significant.

results in the accumulation of cellular DGLA upon, dietary GLA supplementation. DGLA formed within the human neutrophil generally resides in the same phospholipid pools as AA. For example, the largest pools of AA and DGLA within the neutrophil lipids are found in phosphatidylethanolamine (PE) molecular species. Additionally, there are significant increases in the amount of DGLA associated with PE after supplementation with GLA (Chilton-Lopez et al., 1996; Johnson et al., 1997). Thus, the AA/DGLA ratio in PE species markedly decreases after supplementation with GLA. Perhaps more importantly, both DGLA and AA are released from membrane phospholipids, particularly PE, after neutrophil stimulation indicating that DGLA is located in membrane phospholipids that are readily accessible to hydrolysis by phospholipase(s). An altered AA/DGLA ratio has functional implications for immune cells (Table 3).

In contrast to these observations made in several inflammatory cell types, dietary GLA supplementation markedly increases circulating levels of both DGLA and AA in humans, suggesting that ingested GLA is both elongated to DGLA and subsequently desaturated to AA *in vivo* by tissues such as the liver (Barham et al., 2000; Johnson et al., 1997; Surette et al., 2003a). Examination of circulating lipoproteins reveals that DGLA and AA are almost exclusively localized in phospholipid pools, and GLA supplementation further enriches phospholipid pools with both DGLA and AA, but does not cause any appreciable change in other serum lipid classes (Johnson et al., 1997). *In vitro* studies utilizing HepG2 liver cells confirm that these transformed liver cells contain the enzymatic capacities to both elongate GLA to DGLA and then to further desaturate it to form AA (Barham et al., 2000).

Thus, *in vivo* GLA metabolism in humans is extremely complex since all cellular compartments do not metabolize it in a uniform manner due to the differential expression of PUFA metabolizing enzymes. GLA supplementation leads to elevated DGLA and has no effect on AA levels in certain inflammatory cells (e.g. neutrophils), but also increases both DGLA and AA levels in circulating lipids. The biological ramification of AA accumulation in circulation of humans is controversial. Some studies suggest that dietary AA has no influence on immune responses, blood lipids, lipoproteins or health in general (Kelley et al., 1997; Nelson et al., 1997). However, other studies show a strong association between elevated levels of AA and the formation of platelet-aggregating endoperoxides and thromboxanes (Hamberg and Samuelsson, 1974; Hamberg et al., 1975; Smith and Lands, 1972; Willis et al., 1974b). Additionally, high levels of AA in humans have also been shown to result in an increased tendency for the secondary irreversible phase of platelet aggregation (Seyberth et al., 1975).

3.2. Impact of (n-3) PUFAs on GLA metabolism

As mentioned in the Introduction, GLA-enriched supplements have also been given in combination with marine (n-3) long chain-PUFA supplements. These supplementation strategies often provide GLA together with the (n-3) long chain-PUFAs, EPA and DHA. There are three primary rationales for using these combinations. First, addition of (n-3) long chain-PUFAs inhibits the conversion of GLA-derived DGLA to AA. *In vitro* experiments show that EPA blocks *FADS1* activity in cultured HepG2 cells (Barham et al., 2000). Additionally, *in vivo* studies that demonstrate that the addition of fish oil (with EPA and DHA) to GLA-enriched diets prevents the accumulation of serum AA in response to GLA without inhibiting accumulation of DGLA in neutrophils (Barham et al., 2000). Other studies show that inclusion of as little as 0.25 g/d EPA+DHA can block GLA-induced elevations in plasma AA levels (Surette et al., 2003a, 2003b).

Secondly, like GLA alone, supplementation with borage+fish oil combinations inhibit leukotriene generation (Barham et al.,

Table 3
Effect of GLA supplementation on functional endpoints of immune cells.

Reference	design	supplement	dose	duration	cells	Functional Outcomes
Fletcher and Ziboh (1990)	<i>In vivo</i> fatty acid supplementation	Borage oil	3% oil (by weight) in diet	12wk	Guinea pig neutrophil	~25% inhibition of fMLP-stimulated superoxide production (vs control diet; $p < 0.005$) No effect on PMA-stimulated superoxide production (vs control diet)
Iversen et al. (1991)	<i>In vitro</i> fatty acid competition	LA and DGLA conc. curve	0–100 μ M	10 m	Human neutrophil	50 μ M DGLA, 75% \downarrow in LTB ₄ generation 50 μ M LA, 60% \downarrow in LTB ₄ generation
Iversen et al. (1992)	<i>In vitro</i> fatty acid treatment	DGLA conc. curve	0–100 μ M	10 m	Human PBMC ^a	50 μ M DGLA, 60% \downarrow in LTB ₄ generation 50 μ M DGLA, > 7-fold \uparrow in PGE ₁ generation 50 μ M DGLA, significant \uparrow in 15-HETrE generation (> 250 ng/10 mil cells) 50% \downarrow LTB ₄ generation (vs olive oil control), <i>ex vivo</i>
Ziboh and Fletcher (1992)	<i>In vivo</i> fatty acid supplementation	Borage oil	0.48 g GLA/d	6wk	Human neutrophil	50% \downarrow LTB ₄ generation (vs olive oil control), <i>ex vivo</i>
Ziboh and Fletcher (1992)	<i>In vivo</i> fatty acid supplementation	Black currant oil	0.48 g GLA/d	6wk	Human neutrophil	50% \downarrow LTB ₄ generation (vs olive oil control), <i>ex vivo</i>
Chapkin et al. (1988a)	<i>In vitro</i> treatment	15-HETrE	0–30 μ M	1 h	Murine peritoneal macrophage	10 μ M 15-HETrE, 90% \downarrow in LTB ₄ generation
Chilton-Lopez et al. (1996)	<i>In vitro</i> treatment	15-HETrE	0–20 μ M		Human neutrophil	10 μ M 15-HETrE, 75% \downarrow in LTB ₄ generation
Johnson et al. (1997)	<i>In vivo</i> fatty acid supplementation	Borage oil+controlled diet	3 g GLA/day	21d	Human neutrophil	58% \downarrow LTB ₄ generation (vs baseline), <i>ex vivo</i>
Barham et al. (2000)	<i>In vivo</i> fatty acid supplementation	Borage + fish oils + controlled diet	3 g GLA + 3 g EPA/day	21d	Human neutrophil	30% \downarrow LTB ₄ generation (vs baseline), <i>ex vivo</i>
Amagai et al. (2015)	<i>In vitro</i> fatty acid treatment	DGLA conc. curve	0–30 μ M	48 h	RBL-2H3 cells	30 μ M, significant (> 30 ng/ml) PGD ₁ formation
Amagai et al. (2015)	<i>In vivo</i> fatty acid supplementation	DGLA	11% of dietary fatty acids	5wk	NC/Tnd mouse skin	significant \uparrow in PDG ₁ , PDE ₁ , PGD ₂ , 8-HETrE, 15-HETrE (vs control diet)

^a PBMC, peripheral blood mononuclear cells.

2000; Surette et al., 2003a, 2003b) and attenuate the expression of early steps in signal transduction, as well as the expression of genes for pro-inflammatory cytokines (Weaver et al., 2009). Finally, addition of fish oil to GLA supplemented diets enriches cells and tissues with EPA, DPA and DHA and their metabolites. Many of these metabolites have potent anti-inflammatory effects (Ariel and Serhan, 2007; Serhan et al., 2004; Serhan et al., 2002). Consequently, GLA/(n-3) long chain-PUFA combinations theoretically would induce a powerful combination of anti-inflammatory metabolites from DGLA, EPA and DHA.

Botanical oil combinations that contain borage oil, enriched in GLA, and echium oil, (from *Echium plantagineum* L.) enriched in (n-3) PUFAs (ALA and SDA), also markedly increase circulating levels of DGLA and have little impact on circulating AA levels (Arm et al., 2013; Lee et al., 2014). These studies suggest that botanical (n-3) 18C-PUFAs not only enhance the conversion of dietary GLA to DGLA but also inhibit further conversion of that DGLA to AA.

3.3. Differential function and impact of DGLA- and AA-derived eicosanoids

Free (unesterified) DGLA and AA released by phospholipases A₂(s) are substrates for cyclooxygenases (COX) and lipoxygenases (LOX), leading to the synthesis of a variety of eicosanoid products including PGs, TXs, LTs and hydroxyl epoxides. As mentioned above (Section 2.2), there are many eicosanoid derivatives of AA. In particular, the 2-series PGs and TXs, and the 4-series LTs, are by far the most commonly studied and are very well characterized. These lipid mediators tend to exhibit pro-inflammatory activities in numerous cell types and disease states. Additionally, there is emerging scientific literature revealing that free AA and oxidized products of AA can regulate gene expression, and consequently a wide range of cellular activities via cellular and nuclear receptors (Berger et al., 2006; Caputo et al., 2011; Chinetti et al., 2000; Deckelbaum et al., 2006; Jump and Clarke, 1999; Jung et al., 2012; Schmitz and Ecker, 2008; Soberman and Christmas, 2003; Vanden Heuvel, 2012).

Depending on the cell type, DGLA can be metabolized by COX 1/2 (prostaglandin endoperoxide H synthase-1 and -2, *PGHS1/2*) to 1-series PGs, particularly PGE₁, and by 15-lipoxygenase into 15-(S)-hydroxy-8,11,13-eicosatrienoic acid (15-HETrE) (Borgeat et al., 1976). These two metabolites of DGLA have been shown to suppress inflammation, promote vasodilation, lower blood pressure, inhibit smooth muscle cell proliferation, and exert anti-neoplastic activities (Fan et al., 1995; Tabolacci et al., 2010; Zurier, 1991). DGLA supplementation has also been observed to regulate PGD₁ and PGD₂ levels (Amagai et al., 2015) (Table 3).

To better understand the capacity of *PGHS1/2* to synthesize PGE₁ and PGE₂, it is important to understand 1) the cells that contain *PGHS1/2* activities, 2) conditions in which *PGHS1* and *PGHS2* genes are expressed, 3) whether *PGHS1* and *PGHS2* can equally utilize DGLA and AA as substrates for prostaglandin biosynthesis, and 4) the capacity of PGE₁ and PGE₂ to bind E-type prostanoid receptors and induce biological responses. A great deal remains unknown in all of these areas. Levin and colleagues compared the DGLA and AA affinities and reaction rates for *PGHS1* and *PGHS2*. DGLA and AA had similar affinities (Km) and maximal reaction rates (Vmax) for *PGHS2* (Levin et al., 2002). In contrast, AA was metabolized preferentially by *PGHS1*. *PGHS2* (Levin et al., 2002) is thought to be the dominant source for prostaglandin formation during inflammation (Ricciotti and FitzGerald, 2011). Together, these data suggest that both DGLA and AA can be efficiently converted to PGE₁ and PGE₂ during inflammatory responses.

Prostaglandins, including PGE₁ and PGE₂, exert their effects by binding to rhodopsin-like seven transmembrane spanning G

protein-coupled receptors (Ricciotti and FitzGerald, 2011). The prostanoid receptor family has several members including EP1 (E prostanoid receptor 1), EP2, EP3, and EP4 subtypes and while it has been suggested that certain biological properties of PGE₁ are ~20 times stronger than PGE₂ (Fan and Chapkin, 1998), much remains to be learned about the affinity of PGE₁ compared to PGE₂ for PGE receptor subtypes and their subsequent biological activities.

Both PGE₁ and 15-HETrE have been shown to antagonize the synthesis of AA-derived eicosanoids. When exogenously provided, 15-HETrE reduces A23187-stimulated LT generation in a dose-dependent manner in human neutrophils (Chilton-Lopez et al., 1996), peripheral blood mononuclear cells (Iversen et al., 1992) and murine peritoneal macrophages (Chapkin et al., 1988a) by 75–90% (Table 3). The inhibitory effect of 15-HETrE on neutrophil LTB₄ generation is reversible and required only short (5 min) exposure time (Chilton-Lopez et al., 1996). Interestingly, 15-HETrE is a much more potent inhibitor (90% inhibition at 20 μM) of 5-lipoxygenase than 15-lipoxygenase derivatives of AA (15-HETE; 25% at 20 μM), EPA (15-HEPE; 33% at 20 μM) and DHA (17-HODHE; 33% at 20 μM) (Iversen et al., 1992; Ziboh, 1996). PGE₁ appears to also contribute to the inhibitory impact of GLA supplementation on LT synthesis. In mouse dendritic cells, both PGE₁ and PGE₂ suppress LTB₄ production, the latter by an IL-10 dependent mechanism that interferes with the 5-LO activating protein (FLAP) expression (Harizi et al., 2003). Taken together, the isomeric series of the lipid mediators synthesized from AA and DGLA are functionally distinct and typically have opposing actions.

While all putative mechanisms have not been elucidated, the functional consequence of elevated DGLA content in neutrophils is a dramatic reduction in LTB₄ generation in response to stimulation (Chilton-Lopez et al., 1996; Johnson et al., 1997; Ziboh and Fletcher, 1992). Cysteinyl LTs are important in the pathology of asthma and utilize receptors (CysLT₁T and CysLT₂R) localized in bronchial smooth muscle, vascular endothelium and secretory cells (Heise et al., 2000; Lynch et al., 1999). Basophil cysteinyl LT generation is reduced up to 50% in mild asthmatics by supplementation with botanical oil (borage+echium) combinations that contain both GLA and SDA (Arm et al., 2013). The generation of other potent lipid mediators including platelet activating factor is also inhibited as a result of dietary supplementation with GLA (Johnson et al., 1997).

4. The impact of genetic variation in the fatty acid desaturase (*FADS*) gene cluster on AA/DGLA ratios and eicosanoid production

Until recently, the conversion of LA and ALA to AA and DHA, respectively, via the pathway shown in Fig. 1 was thought to be inefficient and uniform for all populations. However, mounting evidence indicates that common genetic and epigenetic variations in close proximity to and within the *FADS* cluster markedly affect the rate of conversion of 18C-PUFAs, including GLA, to LC-PUFAs and thus affecting the amount of circulating and tissue LC-PUFA levels. Specifically, single nucleotide polymorphisms (SNPs) and the methylation status of CpG sites in the *FADS* gene cluster are strongly associated with DGLA, AA and DHA levels in plasma and liver tissues (Chilton et al., 2014; Hoile et al., 2014; Howard et al., 2014; Mathias et al., 2014). As discussed in Section 2, the human *FADS* gene cluster is located on chromosome 11q12–q13.1, comprised of 91.9 kb and has nearly 500 SNPs annotated to this region with exon/intron organization (Fig. 2) (Glaser et al., 2010).

Recent studies from our lab have revealed dramatic population-based differences in the frequency of genetic variations that impact long chain (LC; > 20 carbons) PUFAs and their metabolites




Fig. 2. Plasma AA/DGLA ratios vary by genotype at rs174537 in both African American and European American populations (Adapted from (Mathias et al., 2011)).

(e.g. eicosanoids) (Hester et al., 2014; Mathias et al., 2011; Sergeant et al., 2012). These studies clearly support that associations between genetic variants and PUFA levels are strongly related to ethnicity. For example ~80% of African Americans carry two copies of the alleles associated with increased levels of AA and DHA and reduced levels of DGLA, compared to only ~45% of European Americans. Together these studies raise important questions of whether gene-PUFA interactions induced by a modern western diet are differentially driving the risk of diseases of inflammation in diverse populations, and are these interactions leading to health disparities. They also suggest that “one size fits all” dietary recommendations and supplementation strategies may not be appropriate for all populations or even individuals within specific populations.

Our laboratory has focused on numerous SNPs and epigenetic sites within the *FADS* cluster but particularly the SNP, rs174537. Rs

174,537, located 13–15 kb downstream of *FADS1*, was originally chosen because it was shown in a large GWAS to be the strongest genetic determinant associated circulating plasma AA levels ($p=5.95 \times 10^{-46}$) (Tanaka et al., 2009). Additionally, there are marked frequency differences in genotypes at rs174537 between African-Americans and European-Americans. Our studies, across several cohorts, show that this SNP in particular, is robustly associated with ratios of AA to DGLA and thus the enzymatic efficiency of *FADS1*. Fig. 2 shows the relationships between genotypes at rs174537 and serum AA/DGLA ratios in African American and European American populations (Mathias et al., 2011). These data indicate that there is a greater than 3-fold difference (e.g. African American GG versus European American TT) in the AA/DGLA ratio between all genotypes in both populations and a greater than 2-fold difference between genotypes within either population. African Americans with the GG genotype have a mean ratio of AA/DGLA approaching 7.5–1, with some individuals well over 10–1. Studies have not been performed to determine how genetic variation within the *FADS* cluster impacts AA/DGLA in tissue or inflammatory cell lipids. These observations indicate the critical need for studies that are focused on the impact of gene variations, such as rs174537, on AA/DGLA ratios after supplementation with GLA-enriched oils. However, findings to date suggest that variation within the *FADS* cluster is likely to have significant impact on how individuals respond to GLA supplementation.

5. Discussion and future directions

This review emphasizes that the study of GLA and DGLA metabolism and its relationship to eicosanoid biosynthesis and inflammatory processes is a complex area of research. On the one hand, there are promising studies that suggest that supplementation with GLA and particularly combinations of GLA with (n-3) long chain-PUFAs have great potential to dampen inflammatory processes and improve signs and symptoms of several inflammatory diseases. However, as a whole, this field of study is currently riddled with confusion. Much of the perplexity arises from many of the issues raised in this review including a limited knowledge about how genetic variation affects PUFA supplementation and subsequent metabolism.

A critical question is where does the field go from here? First, even in its current state, we feel the clinical studies indicate that this is an important area of research that should continue to be emphasized. Currently, the clinical effectiveness of a wide variety of supplementation strategies (with fish, flaxseed and GLA-containing oils) is being questioned. For example, recent clinical trials and meta-analyses have challenged the efficacy of supplementation with fish oils containing (n-3) long chain-PUFA (Chen et al., 2011; Filion et al., 2010; Rizos et al., 2012; Zhao et al., 2009). Similarly, a meta-analysis of 27 studies (Pan et al., 2012) showed higher ALA exposure was associated with a moderately lower risk of CVD, but found “high unexplained heterogeneity” that warranted further studies. A central issue in all of these studies is the fact that large, diverse clinical trials inevitably have sizeable subsets of individuals with markedly different circulating and tissue levels of 18C- and LC-PUFAs, and great variability in how individuals respond to PUFA-based supplements. It seems clear that in light of such host-related complexities, study approaches that provide complex (n-6) or (n-3) dietary supplements to diverse groups of individuals using a “one size fits all” model, are unlikely to yield conclusive results.

In contrast, when genetic diversity is taken into consideration and the resultant marked differences in (n-6) to (n-3) LC-PUFA levels and ratios are recognized, then complex (n-6) or (n-3) dietary supplementation strategies can be used to correct critical

diet-gene interactions in a targeted manner for individuals that need them. Moreover, *in vitro*, animal and human studies have demonstrated the benefits of balancing (n-6) and (n-3) metabolic pathways to reduce inflammatory processes, prevent disease and improve human health. It seems that understanding and recognizing individual and population differences provides this field with a great opportunity to optimize the use of PUFA-based supplements (including GLA-enriched supplements) as we move into the era of *individualized medicine*.

Conflict of interest

Floyd H. Chilton is a paid consultant for Eagle Wellness, LLC. This information has been revealed to Wake Forest Baptist Medical Center and is institutionally managed. Other authors have no conflict of interest.

Acknowledgement

This work was supported by NIH P50 AT002782, United States (FHC, PI) and NIH, United States AT008621-01A1 (FHC, PI).

References

- Amagai, Y., et al., 2015. Dihomo- γ -linolenic acid prevents the development of atopic dermatitis through prostaglandin D1 production in NC/Tnd mice. *J. Derm. Sci.* 79, 30–37.
- Andreassi, M., et al., 1997. Efficacy of gamma-linolenic acid in the treatment of patients with atopic dermatitis. *J. Int. Med. Res.* 25, 266–274.
- Ariel, A., Serhan, C.N., 2007. Resolvins and protectins in the termination program of acute inflammation. *Trends Immunol.* 28, 176–183.
- Arm, J., et al., 2013. Impact of botanical oils on polyunsaturated fatty acid metabolism and leukotriene generation in mild asthmatics. *Lipids Health Dis.* 12, 141.
- Bamford, J.T., et al., 2013. Oral evening primrose oil and borage oil for eczema. *Cochrane Database Syst. Rev.* (Art. No. CD004416)
- Barham, J.B., et al., 2000. Addition of eicosapentaenoic acid to γ -linolenic acid-supplemented diets prevents serum arachidonic acid accumulation in humans. *J. Nutr.* 130, 1925–1931.
- Belch, J.J., Hill, A., 2000. Evening primrose oil and borage oil in rheumatologic conditions. *Am. J. Clin. Nutr.* 71, 352s–356s.
- Berger, A., et al., 2006. How dietary arachidonic- and docosahexaenoic- acid rich oils differentially affect the murine hepatic transcriptome. *Lipids Health Dis.* 5, 10.
- Bernert Jr, J.T., Sprecher, H., 1975. Studies to determine the role rates of chain elongation and desaturation play in regulating the unsaturated fatty acid composition of rat liver lipids. *Biochim Biophys. Acta* 398, 354–363.
- Billah, M.M., et al., 1985. Lipoxigenase products of arachidonic acid modulate biosynthesis of platelet-activating factor (1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine) by human neutrophils via phospholipase A2. *J. Biol. Chem.* 260, 6899–6906.
- Blond, J.P., Bézard, J., 1991. Δ 5-desaturation of dihomogammalinolenic acid (20:3 (n-6)) into arachidonic acid (20: 4(n-6)) by rat liver microsomes and incorporation of fatty acids in microsomal phospholipids. *Biochim. Biophys. Acta* 1084, 255–260.
- Borgeat, P., et al., 1976. Transformation of arachidonic acid and homo-gamma-linolenic acid by rabbit polymorphonuclear leukocytes. Monohydroxy acids from novel lipoxygenases. *J. Biol. Chem.* 251, 7816–7820.
- Boyce, J., 2008. Eicosanoids in asthma, allergic inflammation, and host defense. *Curr. Mol. Med.* 8, 335–349.
- Burr, G.O., Burr, M.M., 1929. A new deficiency disease produced by the rigid exclusion of fat from the diet. *J. Biol. Chem.* 82, 345–367.
- Burr, G.O., et al., 1932. On the fatty acids essential in nutrition. III. *J. Biol. Chem.* 97, 1–9.
- Calder, P.C., 2013. Long chain fatty acids and gene expression in inflammation and immunity. *Cur Opin. Clin. Nutr. Metab. Care* 16, 425–433.
- Cameron, M., et al., 2011. Herbal therapy for treating rheumatoid arthritis. *Cochrane Database Syst. Rev.* . <http://dx.doi.org/10.1002/14651858.CD002948.pub2>
- Caputo, M., Zirpoli, H., Torino, G., Tecce, M.F., 2011. Selective regulation of UGT1A1 and SREBP-1c mRNA expression by docosahexaenoic, eicosapentaenoic, and arachidonic acids. *J. Cell. Physiol.* 226, 187–193.
- Chapkin, R.S., Coble, K.J., 1991. Utilization of gammalinolenic acid by mouse peritoneal macrophages. *Biochim Biophys. Acta* 1085, 365–370.
- Chapkin, R.S., et al., 1988a. Ability of 15-hydroxyeicosatrienoic acid (15-OH-20:3) to modulate macrophage arachidonic acid metabolism. *Biochem. Biophys. Res. Comm.* 153, 799–804.
- Chapkin, R.S., et al., 1988b. Inability of murine peritoneal macrophages to convert linoleic acid into arachidonic acid. Evidence of chain elongation. *J. Immunol.* 140, 2350–2355.
- Chapkin, R.S., Ziboh, V.A., 1984. Inability of skin enzyme preparations to bio-synthesize arachidonic acid from linoleic acid. *Biochem. Biophys. Res. Comm.* 124, 784–792.
- Chen, Q., et al., 2011. Effects of omega-3 fatty acid for sudden cardiac death prevention in patients with cardiovascular disease: a contemporary meta-analysis of randomized, controlled trials. *Cardiovas Drugs Ther.* 25, 259–265.
- Chilton-Lopez, T., et al., 1996. Metabolism of gammalinolenic acid in human neutrophils. *J. Immunol.* 156, 2941–2947.
- Chilton, F.H., et al., 2014. Diet-gene interactions and PUFA metabolism: a potential contributor to health disparities and human diseases. *Nutrients* 6, 1993–2022.
- Chinetti, G., et al., 2000. Peroxisome proliferator-activated receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism and inflammation. *Inflam. Res.* 49, 497–505.
- de Bravo, M.M., et al., 1985. Metabolism of gammalinolenic acid by human blood platelet microsomes. *Biochem Int.* 10, 889–896.
- Deckelbaum, R.J., et al., 2006. n-3 Fatty acids and gene expression. *Am. J. Clin. Nutr.* 83, S1520–S1525S.
- Eaton, S.B., 1992. Humans, lipids and evolution. *Lipids* 27, 814–820.
- Fan, Y.-Y., Chapkin, R.S., 1998. Importance of dietary γ -linolenic acid in human health and nutrition. *J. Nutr.* 128, 1411–1414.
- Fan, Y.-Y., et al., 1995. Dietary γ -linolenic acid modulates macrophage-vascular smooth muscle cell interactions: evidence for a macrophage-derived soluble factor that downregulates DNA synthesis in smooth muscle cells. *Arter. Thromb. Vasc. Biol.* 15, 1397–1403.
- Filion, K., et al., 2010. Omega-3 fatty acids in high-risk cardiovascular patients: a meta-analysis of randomized controlled trials. *BMC Cardiovasc Dis.* 10, 24.
- Fiocchi, A., et al., 1994. The efficacy and safety of γ -linolenic acid in the treatment of infantile atopic dermatitis. *J. Int. Med. Res.* 22, 24–32.
- Fletcher, M.P., Ziboh, V.A., 1990. Effects of dietary supplementation with eicosapentaenoic acid or gamma-linolenic acid on neutrophil phospholipid fatty acid composition and activation responses. *Inflammation* 14, 585–597.
- Foolad, N., et al., 2013. Effect of nutrient supplementation on atopic dermatitis in children: A systematic review of probiotics, prebiotics, formula, and fatty acids. *JAMA Derm.* 149, 350–355.
- Foster, R.H., et al., 2010. Borage oil in the treatment of atopic dermatitis. *Nutrition* 26, 708–718.
- Gadek, J.E., et al., 1999. Effect of enteral feeding with eicosapentaenoic acid, gamma-linolenic acid, and antioxidants in patients with acute respiratory distress syndrome. *Crit. Care Med.* 27, 1409–1420.
- Glaser, C., et al., 2010. Role of FADS1 and FADS2 polymorphisms in polyunsaturated fatty acid metabolism. *Metabolism* 59, 993–999.
- Glaser, C., et al., 2011. Genetic variation in polyunsaturated fatty acid metabolism and its potential relevance for human development and health. *Matern. Child. Nutr.* 7, 27–40.
- Hamberg, M., Samuelsson, B., 1974. Prostaglandin endoperoxides. Novel transformations of arachidonic acid in human platelets. *Proc. Nat. Acad. Sci.* 71, 3400–3404.
- Hamberg, M., et al., 1975. Thromboxanes: a new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc. Nat. Acad. Sci.* 72, 2994–2998.
- Harizi, H., et al., 2003. Prostaglandins inhibit 5-lipoxygenase-activating protein expression and leukotriene B4 production from dendritic cells via an IL-10-dependent mechanism. *J. Immunol.* 170, 139–146.
- Heise, C.E., et al., 2000. Characterization of the human cysteinyl leukotriene 2 receptor. *J. Biol. Chem.* 275, 30531–30536.
- Hester, A.G., et al., 2014. Relationship between a common variant in the fatty acid desaturase (FADS) cluster and eicosanoid generation in humans. *J. Biol. Chem.* 289, 22482–22489.
- Hibbeln, J.R., et al., 2006. Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity. *Am. J. Clin. Nutr.* 83, S1483–S1493S.
- Hoile, S.P., et al., 2014. Supplementation with (n-3) long-chain polyunsaturated fatty acids or olive oil in men and women with renal disease induces differential changes in the DNA methylation of FADS2 and ELOVL5 in peripheral blood mononuclear cells. *PLoS One* 9, e109896.
- Horrobin, D.F., 2000. Essential fatty acid metabolism and its modification in atopic eczema. *Am. J. Clin. Nutr.* 71, 367s–372s.
- Howard, T.D., et al., 2014. DNA methylation in an enhancer region of the FADS cluster is associated with FADS activity in human liver. *PLoS One* 9, e97510.
- Hurtado de Catala, G.E., et al., 1992. Arachidonic acid biosynthesis in non-stimulated and adrenocorticotropic-stimulated Sertoli and Leydig cells. *Lipids* 27, 593–598.
- Irazu, C.E., et al., 1993. Delta 5 desaturase activity in rat kidney microsomes. *Mol. Cell. Biochem.* 129, 31–37.
- Iversen, L., et al., 1991. Linoleic acid and dihomogammalinolenic acid inhibit leukotriene B4 formation and stimulate the formation of their 15-lipoxygenase products by human neutrophils *in vitro*. Evidence of formation of anti-inflammatory compounds. *Agents Actions* 33, 286–291.
- Iversen, L., et al., 1992. Effect of dihomogammalinolenic acid and its 15-lipoxygenase metabolite on eicosanoid metabolism by human mononuclear leukocytes *in vitro*: selective inhibition of the 5-lipoxygenase pathway. *Arch. Derm. Res.* 284, 222–226.
- Johnson, M.M., et al., 1997. Dietary supplementation with gamma-linolenic acid

- alters fatty acid content and eicosanoid production in healthy humans. *J. Nutr.* 127, 1435–1444.
- Jump, D.B., Clarke, S.D., 1999. Regulation of gene expression by dietary fat. *Ann. Rev. Nutr.* 19, 63–90.
- Jung, U.J., et al., 2012. Fatty acids regulate endothelial lipase and inflammatory markers in macrophages and in mouse aorta: a role for PPAR γ . *Arter. Thromb. Vasc. Biol.* 32, 2929–2937.
- Kavanagh, K., et al., 2013. Stearidonic and γ -linolenic acids in echium oil improves glucose disposal in insulin resistant monkeys. *Prostaglandin Leukotr. Ess. Fat. Acid.* 89, 39–45.
- Kawamura, A., et al., 2011. Dietary supplementation of gamma-linolenic acid improves skin parameters in subjects with dry skin and mild atopic dermatitis. *J. Oleo Sci.* 60, 597–607.
- Kazani, S., et al., 2014. LTC4 synthase polymorphism modifies efficacy of botanical seed oil combination in asthma. *SpringerPlus* 3, 1–8.
- Kelley, D.S., et al., 1997. Effects of dietary arachidonic acid on human immune response. *Lipids* 32, 449–456.
- Kitz, R., et al., 2006. Impact of early dietary gamma-linolenic acid supplementation on atopic eczema in infancy. *Pedia. Allergy Immunol.* 17, 112–117.
- Kunkel, S., et al., 1981. Suppression of chronic inflammation by evening primrose oil. *Prog. Lipid Res.* 20, 885–888.
- Lagarde, M., et al., 1981. Human platelet PGE1 and dihomogammalinolenic acid. Comparison to PGE2 and arachidonic acid. *Prog. Lipid Res.* 20, 439–443.
- Lee, T.C., et al., 2014. The impact of polyunsaturated fatty acid-based dietary supplements on disease biomarkers in a metabolic syndrome/diabetes population. *Lipids Health Dis.* 13, 196.
- Leventhal, L., et al., 1994. Treatment of rheumatoid arthritis with blackcurrant seed oil. *Br. J. Rheuma.* 33, 847–852.
- Leventhal, L., et al., 1993. Treatment of rheumatoid arthritis with gammalinolenic acid. *Ann. Intern Med.* 119, 867–873.
- Levin, G., et al., 2002. Differential metabolism of dihomogamma-linolenic acid and arachidonic acid by cyclooxygenase-1 and cyclooxygenase-2: implications for cellular synthesis of prostaglandin E1 and prostaglandin E2. *Biochem. J.* 365, 489–496.
- Li, C., et al., 2015. Enteral immunomodulatory diet (omega-3 fatty acid, γ -linolenic acid and antioxidant supplementation) for acute lung injury and acute respiratory distress syndrome: an updated systematic review and meta-analysis. *Nutrients* 7, 5239.
- Lovell, C., et al., 1981. Treatment of atopic eczema with evening primrose oil. *Lancet* 1, 278.
- Luthria, D.L., Sprecher, H., 1994. A comparison of the specific activities of linoleate and arachidonate in liver, heart and kidney phospholipids after feeding rats ethyl linoleate-9,10,12,13-d4. *Biochim. Biophys. Acta* 1213, 1–4.
- Lynch, K.R., et al., 1999. Characterization of the human cysteinyl leukotriene CysLT1 receptor. *Nature* 399, 789–793.
- Macfarlane, G., et al., 2011. Evidence for the efficacy of complementary and alternative medicines in the management of rheumatoid arthritis: a systematic review. *Rheumatol* 50, 1672–1683.
- Mathias, R., et al., 2011. The impact of FADS genetic variants on omega-6 polyunsaturated fatty acid metabolism in African Americans. *BMC Genetics* 12, 50.
- Mathias, R.A., et al., 2014. Genetic variants in the FADS Gene: implications for Dietary Recommendations for Fatty Acid Intake. *Cur Nutr. Rep.* 3, 139–148.
- Morse, N.L., Clough, P.M., 2006. A meta-analysis of randomized, placebo-controlled clinical trials of Efamol[®]; evening primrose oil in atopic eczema. Where do we go here light more recent discoveries. *Cur Pharm. Biotechnol.* 7, 503–524.
- Morse, P.F., et al., 1989. Meta-analysis of placebo-controlled studies of the efficacy of Epogam in the treatment of atopic eczema. Relationship between plasma essential fatty acid changes and clinical response. *Br. J. Derm.* 121, 75–90.
- Navarette, R., et al., 1992. Dietary intake of concentrated gamma-linolenic acid (GLA)-enriched oil suppresses cutaneous level of dihomogamma-linolenic acid (DGLA): possible *in vivo* inhibition of microsomal elongation of GLA to DGLA. *Prostaglandins Leukot. Ess. Fat. Acids* 46, 139–144.
- Nelson, G.J., et al., 1997. The effect of dietary arachidonic acid on platelet function, platelet fatty acid composition, and blood coagulation in humans. *Lipids* 32, 421–425.
- Pan, A., et al., 2012. α -Linolenic acid and risk of cardiovascular disease: a systematic review and meta-analysis. *Am. J. Clin. Nutr.* 96, 1262–1273.
- Pawlosky, R., et al., 1994. Essential fatty acid metabolism in the feline: relationship between liver and brain production of long-chain polyunsaturated fatty acids. *J. Lipid Res.* 35, 2032–2040.
- Pontes-Arruda, et al., 2006. Effects of enteral feeding with eicosapentaenoic acid, γ -linolenic acid, and antioxidants in mechanically ventilated patients with severe sepsis and septic shock. *Crit. Care Med.* 34, 2325–2333.
- Riccioni, G., et al., 2004. Effect of the two different leukotriene receptor antagonists, montelukast and zafirlukast, on quality of life: a 12-week randomized study. *Allergy Asthma Proc.* 25, 445–448.
- Ricciotti, E., FitzGerald, G.A., 2011. Prostaglandins and Inflammation. *Arter. Thromb. Vasc. Biol.* 31, 986–1000.
- Rice, T.W., et al., 2011. Enteral omega-3 fatty acid, γ -linolenic acid, and antioxidant supplementation in acute lung injury. *JAMA* 306, 1574–1581.
- Rizos, E.C., et al., 2012. Association between omega-3 fatty acid supplementation and risk of major cardiovascular disease events: a systematic review and meta-analysis. *JAMA* 308, 1024–1033.
- Schmitz, G., Ecker, J., 2008. The opposing effects of (n-3) and (n-6) fatty acids. *Prog. Lipid Res.* 47, 147–155.
- Sergeant, S., et al., 2012. Differences in arachidonic acid levels and fatty acid desaturase (FADS) gene variants in African Americans and European Americans with diabetes or the metabolic syndrome. *Br. J. Nutr.* 107, 547–555.
- Serhan, C., et al., 2004. Resolvins, docosatrienes, and neuroprotectins, novel omega-3-derived mediators, and their endogenous aspirin-triggered epimers. *Lipids* 39, 1125–1132.
- Serhan, C.N., et al., 2002. Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J. Exp. Med.* 196, 1025–1037.
- Seyberth, H.W., et al., 1975. Increased arachidonate in lipids after administration to man: effects on prostaglandin biosynthesis. *Clin. Pharm. Ther.* 18, 521–529.
- Simon, D., et al., 2014. Gamma-linolenic acid levels correlate with clinical efficacy of evening primrose oil in patients with atopic dermatitis. *Adv. Ther.* 31, 180–188.
- Simopoulos, A.P., 2008. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp. Biol. Med.* 233, 674–688.
- Singer, P., et al., 2006. Benefit of an enteral diet enriched with eicosapentaenoic acid and gamma-linolenic acid in ventilated patients with acute lung injury. *Crit. Care Med.* 34, 1033–1038.
- Smith, W.L., Lands, W.E., 1972. Oxygenation of polyunsaturated fatty acids during prostaglandin biosynthesis by sheep vesicular gland. *Biochemistry* 11, 3276–3285.
- Soberman, R.J., Christmas, P., 2003. The organization and consequences of eicosanoid signaling. *J. Clin. Investig.* 111, 1107–1113.
- Sprecher, H., 1981. Biochemistry of essential fatty acids. *Prog. Lipid Res.* 20, 13–22.
- Surette, M.E., et al., 2003a. Inhibition of leukotriene synthesis, pharmacokinetics, and tolerability of a novel dietary fatty acid formulation in healthy adult subjects. *Clin. Ther.* 25, 948–971.
- Surette, M.E., et al., 2003b. Inhibition of leukotriene biosynthesis by a novel dietary fatty acid formulation in patients with atopic asthma: a randomized, placebo-controlled, parallel-group, prospective trial. *Clin. Ther.* 25, 972–979.
- Surette, M.E., et al., 2008. The impact of a medical food containing gammalinolenic and eicosapentaenoic acids on asthma management and the quality of life of adult asthma patients. *Cur Med. Res. Opin.* 24, 559–567.
- Tabolacci, C., et al., 2010. Similar antineoplastic effects of nimesulide, a selective COX-2 inhibitor, and prostaglandin E1 on B16-F10 murine melanoma cells. *Melanoma Res.* 20, 273–279.
- Tanaka, T., et al., 2009. Genome-wide association study of plasma polyunsaturated fatty acids in the InCHIANTI study. *PLoS Genet.* e1000338.
- Tate, G., et al., 1989. Suppression of acute and chronic inflammation by dietary gamma linolenic acid. *J. Rheuma.* 16, 729–734.
- van Gool, C.J.A.W., et al., 2003. γ -Linolenic acid supplementation for prophylaxis of atopic dermatitis—a randomized controlled trial in infants at high familial risk. *Am. J. Clin. Nutr.* 77, 943–951.
- Van Gool, C.J.A.W., et al., 2004. Oral essential fatty acid supplementation in atopic dermatitis—a meta-analysis of placebo-controlled trials. *Br. J. Derm.* 150, 728–740.
- Vanden Heuvel, J.P., 2012. Nutrigenomics and nutrigenetics of ω 3 polyunsaturated fatty acids. In: Bouchard, C., Ordovas, J.M. (Eds.), *Progress in Molecular Biology and Translational Science*. Academic Press, pp. 75–112.
- Vanderhoeve, J.Y., et al., 1980. Inhibition of leukotriene biosynthesis by the leukocyte product 15-hydroxy-5,8,11,13-eicosatetraenoic acid. *J. Biol. Chem.* 255, 10064–10066.
- Weaver, K.L., et al., 2009. Effect of dietary fatty acids on inflammatory gene expression in healthy humans. *J. Biol. Chem.* 284, 15400–15407.
- Willis, A.L., et al., 1974a. Dihomo-gamma-linolenate suppresses platelet aggregation when administered *in vitro* or *in vivo*. *Prostaglandins* 8, 509–519.
- Willis, A.L., et al., 1974b. An endoperoxide aggregator (Lass), formed in platelets in response to thrombotic stimuli: purification, identification and unique biological significance. *Prostaglandins* 8, 453–507.
- Zhao, Y.-T., et al., 2009. Prevention of sudden cardiac death with omega-3 fatty acids in patients with coronary heart disease: a meta-analysis of randomized controlled trials. *Ann. Med.* 41, 301–310.
- Ziboh, V.A., 1996. The significance of polyunsaturated fatty acids in cutaneous biology. *Lipids* 31, S249–S253.
- Ziboh, V.A., Fletcher, M.P., 1992. Dose-response effects of dietary gamma-linolenic acid-enriched oils on human polymorphonuclear-neutrophil biosynthesis of leukotriene B4. *Am. J. Clin. Nutr.* 55, 39–45.
- Ziboh, V.A., et al., 2000. Metabolism of polyunsaturated fatty acids by skin epidermal enzymes: generation of antiinflammatory and antiproliferative metabolites. *Am. J. Clin. Nutr.* 71, 361s–366s.
- Ziboh, V.A., et al., 2004. Suppression of leukotriene B(4) generation by ex-vivo neutrophils isolated from asthma patients on dietary supplementation with gammalinolenic acid-containing borage oil: possible implication in asthma. *Clin. Devel Immunol.* 11, 13–21.
- Zurier, R., 1991. Role of prostaglandins E in inflammation and immune responses. *Adv. Prostaglandin Thromboxane Leukot. Res.* 21B, 947–953.
- Zurier, R.B., et al., 1996. Gamma-linolenic acid treatment of rheumatoid arthritis. A randomized, placebo-controlled trial. *Arthritis Rheum.* 39, 1808–1817.